

Additional file 1: Measured size classes of the *Alexandrium fundyense* strains Alex2 and Alex5

<i>A. fundyense</i> strain	Treatment / time			25 th percentile	50 th percentile	75 th percentile	
Alex2	Direct grazing	48 h	Control	26.5 µm	30.8 µm	33.6 µm	
			Grazed	24.3 µm	26.5 µm	29.5 µm	
		72 h	Control	25.9 µm	30.2 µm	32.2 µm	
			Grazed	21.3 µm	23.8 µm	26.5 µm	
	Waterborne-cues	48h	Control	25.4 µm	28.3 µm	32.9 µm	
			Fed	23.8 µm	25.9 µm	28.9 µm	
			Starved	24.3 µm	25.9 µm	29.5 µm	
	Alex5	Direct grazing	48 h	Control	21.3 µm	23.3 µm	25.9 µm
				Grazed	20.4 µm	22.3 µm	23.8 µm
72 h			Control	21.3 µm	23.8 µm	26.5 µm	
			Grazed	20.4 µm	22.3 µm	24.3 µm	
Waterborne-cues		48h	Control	21.3 µm	23.8 µm	27.1 µm	
			Fed	20.9 µm	22.3 µm	24.3 µm	
			Starved	20.9 µm	22.8 µm	24.8 µm	