

Creation of synthetic data

Additional file 3 for “Quantifying Differences in Cell Line Population Dynamics Using CellPD”

Authors: Edwin F. Juarez^{1,2}, Roy Lau¹, Samuel H. Friedman¹, Ahmadreza Ghaffarizadeh¹, Edmond Jonckheere², David B. Agus¹, Shannon M. Mumenthaler¹, and Paul Macklin¹.

Authors' affiliations:

1: Lawrence J. Ellison Institute for Transformative Medicine, University of Southern California.
2: Department of Electrical Engineering, Viterbi School of Engineering, University of Southern California.

Contact information: Paul Macklin: Paul.Macklin@usc.edu
Edwin F. Juarez: juarezro@usc.edu

In order to test the high-throughput capabilities of CellPD we generated a synthetic dataset. The goal of this dataset was to imitate the output file of a high content screening microscope. We simulated a (basic) drug screening experiment testing two compounds. We predefine the seeding density, birth rate, death rate, clearance rate, and sensitivity of a cell line for the mathematical model shown below:

$$\frac{dL}{dt} = \text{birth_rate}(1 - L/L_{max})(1 - \text{sensitivity}_1 \times \text{cytostaticity} \times \text{concentration})L - \text{death_rate}(1 + \text{sensitivity}_2 \times \text{citotoxicity} \times \text{concentration})L$$

$$\frac{dD}{dt} = \text{death_rate}(1 + \text{sensitivity}_2 \times \text{citotoxicity} \times \text{concentration})L - \text{clearance_rate} \times D$$

$$L(0) = \text{seeding_cells_live}$$

$$D(0) = \text{seeding_cells_dead}$$

$$\text{birth_rate} = 5.5\text{e-}2 \text{ h}^{-1}$$

$$\text{death_rate} = 5.0\text{e-}3 \text{ h}^{-1}$$

$$\text{clearance_rate} = 2.0\text{e-}3 \text{ h}^{-1}$$

$$\text{carrying_capacity} = 1.0\text{e}9 \text{ cells}$$

$$\text{seeding_cells_live} = 1.0\text{e}3 \text{ cells}$$


$$\text{seeding_cells_dead} = 1.0\text{e}1 \text{ cells}$$

$$\text{sensitivity}_1 = 1.0$$

$$\text{sensitivity}_2 = 1.0$$

$$\text{concentration} = 0, 0.2, 0.4, 0.6, 0.8$$

To simulate a cytostatic drug, we set $\text{cytostaticity} = 1.0$ and $\text{citotoxicity} = 0.20$, similarly, to simulate a cytotoxic drug, we set $\text{cytostaticity} = 0.20$ and $\text{citotoxicity} = 1.0$. We simulate three biological replicates for each drug concentration. In order to simulate biological heterogeneity and measurement error, we add a Gaussian noise of 15% to each biological replicate.



Drug Name	Concentration	True birth rate	Estimate	Difference	Percent Error
Drug A	0	0.055	0.055	0	0
Drug A	0.2	0.044	0.0452	0.0012	2.727
Drug A	0.4	0.033	0.0328	0.0002	0.606
Drug A	0.6	0.022	0.0221	0.0001	0.455
Drug A	0.8	0.011	0.0112	0.0002	1.818
Drug B	0	0.055	0.056	0.001	1.818
Drug B	0.2	0.0528	0.0579	0.0051	9.659
Drug B	0.4	0.0506	0.0507	0.0001	0.197
Drug B	0.6	0.0484	0.05	0.0016	3.305
Drug B	0.8	0.0462	0.0486	0.0024	5.195
				Mean percent error:	2.578

Table S9-2: Death rate estimate error					
Drug Name	Concentration	True death rate	Estimate	Difference	Percent Error
Drug A	0	0.005	0.005	0	0
Drug A	0.2	0.0052	0.00559	0.00039	7.5
Drug A	0.4	0.0054	0.00592	0.00052	9.630
Drug A	0.6	0.0056	0.00565	5.00E-05	0.893
Drug A	0.8	0.0058	0.00566	0.00014	2.414
Drug B	0	0.005	0.00564	0.00064	12.8
Drug B	0.2	0.006	0.0078	0.0018	30
Drug B	0.4	0.007	0.00692	8.00E-05	1.143
Drug B	0.6	0.008	0.00799	1.00E-05	0.125
Drug B	0.8	0.009	0.00924	0.00024	2.667
				Mean percent error:	6.717

Table S9-3: Death rate estimate error					
Drug Name	Concentration	True net growth rate	Estimate	Difference	Percent Error
Drug A	0	0.05	0.05	0	0
Drug A	0.2	0.0388	0.03961	0.00081	2.088
Drug A	0.4	0.0276	0.02688	0.00072	2.609
Drug A	0.6	0.0164	0.01645	5.00E-05	0.305
Drug A	0.8	0.0052	0.00554	0.00034	6.538
Drug B	0	0.05	0.05036	0.00036	0.72
Drug B	0.2	0.0468	0.0501	0.0033	7.051
Drug B	0.4	0.0436	0.04378	0.00018	0.413
Drug B	0.6	0.0404	0.04201	0.00161	3.985
Drug B	0.8	0.0372	0.03936	0.00216	5.806
				Mean percent error:	2.952